



Inducible-costimulator-mediated suppression of human immunodeficiency virus type 1 replication in CD4⁺ T lymphocytes

Xin Zhou,^a Makoto Kubo,^a Hironori Nishitsuji,^a Kiyoshi Kurihara,^a Tamako Ikeda,^a Takashi Ohashi,^a Miyuki Azuma,^b Takao Masuda,^a and Mari Kannagi^{a,*}

^aDepartment of Immunotherapeutics, Graduate School, Tokyo Medical and Dental University, Tokyo 113-8519, Japan

^bDepartment of Molecular Immunology, Graduate School, Tokyo Medical and Dental University, Tokyo 113-8519, Japan

Received 8 January 2004; returned to author for revision 24 March 2004; accepted 28 April 2004

Available online 19 June 2004

Abstract

We investigated the effects of signaling through CD28 family molecules on human immunodeficiency virus type 1 (HIV-1) replication in vitro. A monoclonal antibody (mAb) specific for inducible costimulator (ICOS) suppressed both X4 and R5 HIV-1 replication in CD4⁺ peripheral blood mononuclear cells (PBMC). This suppression was not attributable to reduced cell growth or viability. CD28 mAb showed variable effects and also suppressed HIV-1 replication when immobilized. Replication of pseudotype viruses with HIV-1-but not with vesicular stomatitis virus G-envelope was efficiently suppressed in CD4⁺ PBMC treated with ICOS or CD28 mAbs. However, CD4, CXCR4, and CCR5 expression on the surface was not down-regulated. Moreover, HIV-1 replication in CD4⁺ PBMC was suppressed by a soluble form of human B7-H2, a ligand of ICOS, but was enhanced by soluble B7-1, a ligand for CD28. These findings suggest that natural or artificial ligands for ICOS potentially suppress HIV-1 replication mainly at the entry stages.

© 2004 Elsevier Inc. All rights reserved.

Keywords: HIV-1 suppression; Immune regulation; Costimulatory molecule; ICOS; CD28; T cell activation

Introduction

HIV-1 proliferates extensively and creates viremia in the acute phase of HIV-1 infection. Following the acute phase, the peripheral HIV-1 load declines to low levels that are continued during the asymptomatic phase of several years (Ho et al., 1989). Even during the asymptomatic period, viral replication is maintained at a low level (Pantaleo et al., 1993; Piatak et al., 1993). HIV-1 production in vivo is performed by both continuous de novo infections to uninfected short-lived cells, and viral persistence in long-life cells (Ho et al., 1995; Perelson et al., 1996; Wei et al., 1995). Individuals who receive long-term highly active antiretroviral therapy retain replication-competent HIV-1 predominantly in the memory T cell fractions, suggesting that these cells may be the main reservoir of persistently infecting HIV-1 (Finzi et al., 1997; Wong et al., 1997).

The mechanisms regulating HIV-1 replication in vivo are not fully understood. HIV-1-specific cytotoxic T cells (CTL)

are thought to kill HIV-1-infected cells, although Nef-mediated down-regulation of MHC-I may limit CTL function (Walker et al., 1987). CD8⁺ cell-mediated unidentified soluble and cellular molecules may also contribute to the suppression of viral replication (Kannagi et al., 1990; Walker et al., 1986). Neutralizing antibodies might have some effect on limiting the spread of HIV-1. Besides these host defense mechanisms, the status of T cell activation may also affect HIV-1 replication by altering the expression of HIV-1 receptors and cytokines, as well as the status of intracellular molecules responsible for nuclear transport or transcription.

T cell activation is regulated by signals through a T cell receptor/CD3 complex and various costimulatory molecules that interact with their ligands on antigen-presenting cells (APC) (Lenschow et al., 1996). The first well-characterized costimulatory molecule was CD28 (Hara et al., 1985), which mediates positive signals in T cells following interactions with B7-1/CD80 or B7-2/CD86 (Linsley et al., 1990). Inhibition of CD28 receptors prevented CD3-mediated HIV-1 replication, indicating that CD28 signals are crucial in HIV-1 replication (Diegel et al., 1993). In contrast, CD28 may also transmit signals inhibiting HIV-1 replication when ligated (Levine et al., 1996).

* Corresponding author. Fax: +81-3-5803-0235.

E-mail address: kann.impt@tmd.ac.jp (M. Kannagi).

Several other members of the CD28 family molecules have been identified including cytotoxic T lymphocyte antigen 4 (CTLA-4) (Brunet et al., 1987), inducible costimulator (ICOS)/activation-inducible lymphocyte immunomediatory molecule (AILIM)/H4 (Buonfiglio et al., 2000; Hutloff et al., 1999; Tamatani et al., 2000), and programmed death gene 1 (PD-1) (Nishimura and Honjo, 2001). CTLA-4 is an inducible molecule homologous to CD28 and also binds B7-1/CD80 and B7-2/CD86, but transmits negative signals to T cells (Azuma et al., 1993; Krummel and Allison, 1995; Walunas et al., 1994). ICOS transmits positive signals following interactions with the ligand B7-H2/B7 homologous protein (B7h)/B7-related protein 1 (B7RP-1) (Swallow et al., 1999; Yoshinaga et al., 1999). Stimulation of CD28 results in activation of IL-2, while stimulation of ICOS induces the production of other cytokines such as IL-4, IL-5, IL-10, IL-13, IFN- γ , TNF- α , and GM-CSF (Beier et al., 2000; Coyle et al., 2000; Hutloff et al., 1999). PD-1 binds PD-L1 (B7-H1) and PD-L2 (B7-DC) (Dong et al., 1999; Freeman et al., 2000; Latchman et al., 2001; Tseng et al., 2001), and inhibits TCR-mediated proliferation of T cells and cytokine production. CD28 is constitutively expressed by T cells, while other molecules are induced upon activation (Sharpe and Freeman, 2002).

B7 family molecules are expressed on monocytes, dendritic cells, and B cells, and sometimes on T cells, epithelial cells, and endothelial cells (Frauworth and Thompson, 2002; Sharpe and Freeman, 2002). Although B7-1/B7-2 are mainly expressed on lymphoid organs, B7h, PD-L1, and PD-L2 are also found on nonlymphoid organs such as the heart, lung, and kidney (Latchman et al., 2001; Lenschow et al., 1996; Sharpe and Freeman, 2002; Swallow et al., 1999). Thus, interactions between molecules from the CD28-B7 family may play an important role not only in priming T cells, but also in regulating activated T cells in both lymphoid and nonlymphoid organs in vivo.

In the present study, we investigate effects of signals through CD28 family molecules on HIV-1 replication, and showed that ICOS monoclonal antibody (mAb) mediate suppression of HIV-1 replication in PHA-stimulated CD4⁺ T cells in vitro mainly at the early stages of HIV-1 replication. Furthermore, a soluble form of B7-H2 (B7h) also mediated signals suppressing HIV-1 replication, suggesting that natural or artificial ligands for ICOS might be applicable as a biological strategy to control HIV-1 infection.

Results

The effects of mouse monoclonal antibodies (mAbs) for the CD28 family molecules on HIV-1 replication

To examine the effect of activation signals through the CD28 family molecules on HIV-1 replication, phytohemagglutinin (PHA)-stimulated CD4⁺ peripheral blood mononuclear cells (PBMC) were infected by the X4 strain HIV-1 NL4-3, and cultured in the presence of mAbs for various CD28 family molecules at concentrations ranging from 0.5 to 10 μ g/ml. HIV-1 replication was monitored by measuring the levels of HIV-1 p24 in the culture supernatant after 4 days of culture. The results of four experiments using PBMC isolated from different donors are summarized in Table 1. Although there was individual variability, SA12, a mAb to ICOS, significantly suppressed HIV-1 replication. The effects of the anti-CD28 mAb TN228 varied widely, enhancing viral replication in some samples, while suppressing it in others. MAbs to CTLA-4 and PD-1 did not markedly affect HIV-1 replication. Flow cytometric analysis indicated that PHA-stimulated CD4⁺ PBMC expressed significant levels of CD28 and ICOS, but low or undetectable levels of PD-1 and CTLA-4 (Fig. 1). We therefore investigated the effects of ICOS and CD28 signals on HIV-1 replication thereafter.

Table 1
Effects of mAbs to various CD28 family molecules on HIV-1 replication

| | Concentration of mAbs (μ g/ml) | HIV-1 p24 production (pg/ml) in CD4 ⁺ PBMC culture in the presence of | | | | | |
|--------|-------------------------------------|--|------------------|-------------------|------------------|------------------|------------------|
| | | Medium | TN228 anti-CD28 | MIH8 anti-CTLA-4 | SA12 anti-ICOS | MIH4 anti-PD-1 | Mouse IgG1 |
| Exp. 1 | 0 | 130.1 \pm 6.1 | | | | | |
| | 10 | | 147.1 \pm 32.0 | 158.0 \pm 27.0 | 100.0 \pm 12.1 | 147.3 \pm 79.3 | 135.0 \pm 16.1 |
| Exp. 2 | 0 | 152.8 \pm 9.7 | | | | | |
| | 1 | | 123.5 \pm 11.2 | 119.0 \pm 12.9* | 129.1 \pm 9.6 | 134.8 \pm 20.4 | 136.0 \pm 9.6 |
| | 2.5 | | 75.9 \pm 17.5* | 149.6 \pm 13.7 | 58.3 \pm 5.5* | 136.4 \pm 6.2 | 140.5 \pm 8.9 |
| Exp. 3 | 0 | 77.2 \pm 13.8 | | | | | |
| | 10 | | 133.0 \pm 8.6* | 70.2 \pm 20.0 | 39.6 \pm 7.4* | 79.4 \pm 8.3 | 82.0 \pm 11.0 |
| Exp. 4 | 0 | 168.8 \pm 20.6 | | | | | |
| | 0.5 | | 183.7 \pm 14.0 | 138.4 \pm 14.5 | 78.5 \pm 4.4* | 146.2 \pm 14.9 | 151.5 \pm 38.2 |
| | 5 | | 199.2 \pm 59.0 | 149.7 \pm 10.5 | 49.9 \pm 17.3* | 155.1 \pm 72.3 | 141.2 \pm 34.6 |

PHA-stimulated CD4⁺ PBMC cells were infected with the X4 HIV-1 strain NL4-3 for 2 h and cultured either without or with the indicated mAbs at the indicated concentration in 96-well round-bottomed plates. HIV-1 p24 in the supernatant was measured by ELISA after 4 days. The results of four independent experiments using PBMC from different donors are shown. The values indicate the mean \pm SD from triplicate determinations.

* P < 0.05.




Fig. 1. Cell surface expression of CD28 family molecules on PHA-stimulated CD4⁺ PBMC. PHA-stimulated CD4⁺ PBMC cultured in the presence of IL-2 for 6 days were stained with mAbs to CD28, CTLA-4, ICOS, or PD-1 (open histogram), and subsequently stained with FITC-labeled second antibody. Closed histogram indicates cells stained with second antibody alone.

Inhibition of X4 and R5 HIV-1 replication by immobilized ICOS and CD28 mAbs

It has previously been shown that immobilized CD28 mAbs suppress HIV-1 replication in PBMC (Levine et al., 1996). We then assessed the effects of immobilized ICOS and CD28 mAbs on X4 and R5 HIV-1 replication. As shown in Fig. 2, immobilized CD28 and ICOS mAbs significantly inhibited replication of both X4 HIV-1 NL4-3 and R5 HIV-1 JR-CSF in CD4⁺ PBMC. The inhibitory effect of immobilized ICOS and CD28 mAbs on HIV-1 replication was observed in repetitive experiments. Thus, ligation of ICOS or CD28 mediated efficient inhibitory signals in HIV-1 replication in CD4⁺ T cells regardless of X4 or R5 HIV-1 strains.

Immobilized ICOS and CD28 mAbs do not suppress cell proliferation or viability

We next examined whether ICOS- or CD28-mediated inhibition of viral replication is associated with reduced proliferation or viability of CD4⁺ T cells. Fig. 3A shows HIV-1 replication in PHA-stimulated CD4⁺ T cells cultured in either the presence of immobilized ICOS or CD28 mAbs at concentrations ranging from 0.0125 to 5 μ g/ml for 4 days. HIV-1 replication was dose-dependently suppressed by immobilized ICOS or CD28 mAbs, except at the highest concentration of CD28 mAb, which showed some irregularity. Simultaneously prepared uninfected CD4⁺ T cells from the same donor underwent a [³H]thymidine uptake

assay after 4 days of culture in the presence of similar concentrations of immobilized mAbs. As shown in Fig. 3B, in contrast to HIV-1 replication, proliferation of CD4⁺ T cells was positively correlated with the presence of either immobilized ICOS or CD28 mAbs in a dose-dependent manner. As a control, immobilized mouse IgG1 did not affect either HIV-1 replication or cell proliferation. To assess whether cell death occurred, we daily monitored the levels of lactate dehydrogenase (LDH) released into the supernatant from CD4⁺ T cells cultured with immobilized mAbs. As shown in Fig. 3C, the amount of LDH did not differ among these CD4⁺ T cell cultures during 3 days of culture. These results indicated that the inhibition of HIV-1 replication by immobilized ICOS or CD28 mAbs was not due to inhibition of cell proliferation or viability.

ICOS and CD28 mediate inhibition of the early stages of a HIV-1 replication cycle

To determine at which steps of the life cycle of HIV-1 the antiviral effect of ICOS or CD28 mAbs occurred, we monitored HIV-1 cDNA synthesis after infection. PHA-stimulated CD4⁺ T cells infected with HIV-1 NL4-3 for 2 h were cultured with immobilized ICOS or CD28 mAbs, and 10⁶ cells in each well were harvested for DNA analysis 2, 24, and 48 h after infection (Fig. 4). The reverse transcripts were amplified by PCR using specific primer pairs for HIV-1 LTR R and gag regions (R/gag) (Masuda et al., 1995; Zack et al., 1990). Twenty-four hours after infection, the amount of viral cDNA (R/gag) was considerably less in




Fig. 2. Immobilized mAbs for ICOS (SA12) and CD28 (TN228) inhibited HIV-1 replication. CD4⁺ PBMC cultured for 4–6 days after PHA stimulation were infected with X4 HIV-1 NL4-3 (A) or R5 HIV-1 JR-CSF (B) for 2 h, and cultured in medium alone or in the plates precoated with 5 μ g/ml antibodies indicated. After 4 days, HIV-1 p24 levels in the supernatant were measured by ELISA. The results indicate the mean \pm SD from triplicate determinations. Similar results were obtained in an independent experiment. * P < 0.05.

ICOS mAb-treated CD4⁺ T cells compared with the control cells not treated with mAbs. Although to a lesser degree than ICOS mAb, CD28 mAb also inhibited cDNA synthesis of HIV-1. Similar inhibition of viral cDNA synthesis by these mAbs was also observed at 14 h following HIV-1 infection in an independent experiment (data not shown). These results suggest that immobilized ICOS and CD28

mAbs prevent HIV-1 infection during the steps before or at reverse transcription in the HIV-1 life cycle.

The effects of immobilized ICOS and CD28 mAbs on pseudotype HIV-1 expression

We next investigated whether ICOS or CD28 mAbs could suppress the entry steps of HIV-1 replication by using pseudotype HIV-1 (Masuda et al., 1995; Planelles et al., 1995) consisting of an envelope-defective HIV-1 expressing




Fig. 3. Stimulation of ICOS or CD28 suppressed HIV-1 replication but not the proliferation or viability of CD4⁺ T cells. (A) PHA-stimulated purified CD4⁺ T cells infected with HIV-1 NL4-3 were cultured on plates precoated with antibodies at the indicated concentration of 5 μ g/ml (■), 0.25 μ g/ml (▨), and 0.0125 μ g/ml (□), or in the medium alone (□) for 4 days, and HIV-1 p24 levels in the supernatant was measured by ELISA. (B) Uninfected PHA-stimulated purified CD4⁺ T cells from the same donor as A were simultaneously cultured in plates similarly precoated with antibodies for 4 days, and the incorporation of [³H]thymidine (Tdr) into the cells during the last 16 h was measured as described in Materials and methods. The results indicate the mean \pm SD from triplicate determinations. (C) Cell death of PHA-stimulated purified CD4⁺ T cells incubated on plates precoated with antibodies at the indicated concentrations was determined daily from the level of LDH in the supernatant over 3 days. The results are shown for 24 h after the initiation of culture. Similar results were obtained in the second and third days.




Fig. 4. Analysis of HIV-1 DNA synthesis in NL4-3-infected CD4⁺ PBMC treated with immobilized ICOS mAb. CD4⁺ PBMC cultured for 4 days after PHA stimulation were infected with DNase-treated HIV-1 NL4-3 for 2 h and cultured in 12-well flat-bottomed plates (10^6 cells/well) precoated with ICOS or CD28 mAbs (5 μ g/ml) or medium alone. Total DNA was extracted from the cells harvested at the indicated periods after infection and underwent PCR analysis with the primer pairs for HIV-1 R/gag region. For HIV-1 DNA standards, 250–5000 copies of linearized HIV-1 JR-CSF DNA were amplified in parallel. Amplified products were resolved on a 2% agarose gel and visualized by Syber-Green staining (FMC Bioproduct). HIV-1 treated at 65 °C for 30 min before inoculation was used as a heat-inactivated control (HI).

luciferase and either HIV-1 LAI envelope (Poon et al., 1998) or VSV-G envelope (Yee et al., 1994). The pseudotype viruses with HIV-1 envelope enter into cells by envelope-mediated membrane fusion, whereas the pseudotype viruses with VSV-G envelope use endocytotic mechanisms. Fig. 5 shows luciferase activities of PHA-stimulated CD4⁺ PBMC that were infected with pseudotype HIV-1 and cultured in the presence of immobilized ICOS or CD28 mAbs. Luciferase expression in the CD4⁺ PBMC infected with VSV-G envelope pseudotype virus was increased by CD28 mAbs and unchanged by ICOS mAbs (Fig. 5A). By contrast, expression of HIV-1 envelope pseudotype virus was significantly inhibited by both ICOS and CD28 mAbs (Fig. 5B). These results indicate that immobilized ICOS and CD28 mAbs mainly affected the entry stages of HIV-1 replication.

ICOS or CD28 stimulation did not reduce HIV-1 receptor expression on CD4⁺ T cells

We next assessed whether pretreatment with immobilized ICOS or CD28 mAbs reduces the expression of HIV-1-specific receptors on CD4⁺ T cells. Following 24 h of preincubation with immobilized ICOS or CD28 mAbs, the expression of CD4, CXCR4, and CCR5 on PHA-stimulated CD4⁺ cells was analyzed by flow cytometry. The purity of the CD4⁺ T cells was approximately 93%. The results are shown in Fig. 6. The expression of CCR5 was significantly upregulated in cultures pretreated with CD28 or ICOS mAbs. In these cultures, increased CCR5 expression persisted during the first 4 days after stimulation at least. In contrast, the levels of CD4 and CXCR4 expression were not altered by these mAbs. Thus, ICOS- or CD28-mediated suppression of both X4 and R5 HIV-1 replication could not be attributed to a reduction in HIV-1-specific receptor expression on the cell surface.




Fig. 5. Effects of ICOS or CD28 stimulation on pseudotype HIV-1 gene expression. CD4⁺ PBMC cultured for 4 days after PHA stimulation were incubated overnight in plates coated with immobilized ICOS or CD28 mAbs or control mouse IgG (1 μ g/ml) (10^6 /well in 24-well plates), then infected with luciferase-expressing pseudotype HIV-1 possessing a VSV-G envelope (A) or a HIV-1 LAI envelope (B) for 2 h. The cells were further cultured for 48 h on plates coated with the same antibodies as above, and luciferase activity was measured. The results show the mean \pm SD from duplicate determinations. * P < 0.05.




Fig. 6. Effects of ICOS or CD28 stimulation on HIV-1 receptor expression by CD4⁺ T cells. PHA-stimulated CD4⁺ PBMC were treated either with immobilized mAbs to ICOS (A) or CD28 (B) or control mouse IgG1 (C) (5 μ g/ml) in 24-well flat-bottomed plates, and the expression of CXCR4 and CCR5 receptors was examined by FACS (open histogram) after 24 h of culture. Closed histogram indicates control cells stained with the second antibody alone. Similar results were obtained in an independent experiment cultured for 3 days.

Effect of ICOS or CD28 stimulation on cytokine production

We next examined whether ICOS and CD28 mAbs altered the production of cytokines and chemokines that might affect HIV-1 replication. In most of the experiments above, we added ICOS or CD28 stimulation to CD4⁺ PBMC that had been cultured for several days after PHA stimulation. In this condition, the cells stimulated with PHA alone produced detectable levels of IL-5, IFN- γ , MIP-1 α , and RANTES, and very low levels of IL-10 (Table 2). Among them, IL-5 production was further enhanced by CD28-stimulation, but other cytokines were not significantly altered by ICOS or CD28 mAbs.

We also examined cytokine production by CD4⁺ PBMC stimulated with ICOS or CD28 mAbs immediately after PHA stimulation. In this condition, the cells with PHA stimulation alone produced significant levels of IL-5, IL-10, TNF α , IFN γ , RANTES, and MIP-1 α . Stimulation of ICOS and CD28 enhanced IL-5 and IL-10, but reduced TNF α and IFN γ production. RANTES and MIP-1 α were not significantly affected by ICOS and CD28 mAbs. SDF-1 was not detectable in any culture tested.

We also confirmed that the expression of pseudotype virus with HIV-1 envelope was significantly inhibited in the cells that were stimulated with ICOS or CD28

mAbs immediately after PHA stimulation, (Fig. 7A), whereas similar treatment did not affect cell proliferation (Fig. 7B).

Table 2

Effect of ICOS and CD28 stimulation on cytokine production (pg/ml) from PHA-stimulated CD4⁺ PBMC

| Cytokines | Early stimulation with | | | Late stimulation with | | |
|----------------|------------------------|--------|--------|-----------------------|-------|--------|
| | Mouse IgG1 | SA12 | TN228 | Mouse IgG1 | SA12 | TN228 |
| IL-4 | N.D. | 4.4 | 2.3 | N.D. | N.D. | 1.3 |
| IL-5 | 621.3 | 1785.1 | 1514.9 | 679.6 | 773.6 | 1053.4 |
| IL-10 | 171.6 | 274.7 | 246.8 | 8.9 | 11.2 | 13 |
| TNF- α | 1599.2 | 1075.7 | 942.4 | 1.8 | 2.4 | 2.1 |
| IFN- γ | 13814.5 | 2913.4 | 3858.3 | 338.5 | 356.6 | 419 |
| MIP-1 α | 1512.7 | 1629.9 | 1554.5 | 379.1 | 356.7 | 394 |
| RANTES | 3524.4 | 3565.9 | 3177.8 | 426.7 | 400 | 429.6 |
| SDF-1 α | N.D. | N.D. | N.D. | N.D. | N.D. | N.D. |

CD4⁺ PBMC were placed into plates that were precoated with mouse IgG1, or SA12 (anti-ICOS) or TN228 (anti-CD28) mAbs at a concentration of 5 μ g/ml, immediately (early stimulation) or 4 days (late stimulation) after 30 min of PHA stimulation, and cultured in the RPMI medium containing IL-2. Cytokines and chemokines in the supernatants were measured by using either Cytometric Bead Array Kit (IL-4, IL-5, IL-10, TNF- α) or ELISA (IFN- γ , MIP-1 α , RANTES, SDF-1 α) 3 days after mAb stimulations.




Fig. 7. Effects of ICOS or CD28 stimulation immediately after PHA stimulation. CD4⁺ PBMC were stimulated with PHA for 30 min, washed, and cultured on plates precoated with immobilized ICOS or CD28 mAbs or control mouse IgG (1 μ g/ml) in the presence of IL-2. HIV-1 expression (A) and proliferation (B) of these cells were examined. (A) Two days after stimulation, the cells were infected with pseudotype virus with a HIV-1 envelope, and further cultured for 48 h in the mAb-coated wells, and then luciferase activity was measured. * $P < 0.05$. (B) The proliferation of the cells was measured by the [³H]thymidine incorporation method after a 3-day culture without infection. The results show the mean \pm SD from duplicate (A) or quadruplicate (B) determinations.

Leflunomide reversed the inhibition of HIV-1 replication mediated by immobilized ICOS or CD28 mAbs

The intracellular transduction pathway underlying ICOS signaling is not fully understood. Consequently, we investigated the effects of inhibitors for NF- κ B (Leflunomide), MEK (PD98059), and p38 MAP kinase (SB202190) on ICOS- or CD28-induced inhibition of HIV-1 replication. As shown in Fig. 8, Leflunomide significantly attenuated CD28-induced inhibition of HIV-1 replication in CD4⁺ T cells. Although Leflunomide alone suppressed HIV-1 replication, the coexistence of immobilized CD28 mAb resulted in overwhelming viral replication. A low level of enhancement in HIV-1 replication was also observed in ICOS mAb-treated cells in the presence of Leflunomide, although it was not statistically significant. In the presence

of PD98059, CD28 but not ICOS mAb significantly enhanced HIV-1 replication. SB202190 marginally altered the effects of ICOS- or CD28-mAbs. These results suggested that CD28 and ICOS in a lesser degree mediated positive signals in HIV replication besides inhibitory signals, and that the inhibitory signals were suppressed by inhibiting NF- κ B and/or MEK pathways.

Natural ligands for ICOS-inhibited HIV-1 replication

Finally, we assessed whether the natural ligands of ICOS or CD28 affect HIV-1 replication. Recombinant chimeric proteins of various B7-CD28 family molecules and the Fc region of human IgG were added to CD4⁺ T cell cultures




Fig. 8. The effects of Leflunomide on ICOS- or CD28-induced inhibition of HIV-1 replication. (A) PHA-stimulated CD4⁺ PBMC infected with HIV-1 NL4-3 were cultured in the absence (\blacklozenge) or presence of various concentrations of Leflunomide (NF- κ B-inhibitor) (\blacksquare), PD98059 (MEK-inhibitor) (\blacktriangle), or SB202190 (p38 MAP kinase inhibitor) (\times), for 4 days, and HIV-1 p24 levels in the supernatant were measured. (B) PHA-stimulated CD4⁺ PBMC infected with HIV-1 NL4-3 were cultured either without (\square) or with immobilized ICOS (\boxtimes) or CD28 (\boxdot) mAbs (5 μ g/ml) in 48-well flat-bottomed plates in either the absence or presence of 20 μ M Leflunomide, PD98059, or 10 μ M of SB202190. HIV-1 p24 levels in the supernatant were measured by ELISA 4 days after infection. The results indicate the mean \pm SD from triplicate determinations. Similar results were obtained in an independent experiment. * $P < 0.05$.

Table 3
Effects of various soluble B7-CD28 family molecules on HIV-1 replication

| Reagents | HIV-1 p24 (pg/ml) in the supernatant of HIV-1-infected CD4+ PBMC culture | |
|----------------------|--|--------------|
| | Exp. 1 | Exp. 2 |
| Medium | 77.2 ± 13.8 | 168.8 ± 20.6 |
| Mouse IgG1 | 82.0 ± 11.0 | N.T. |
| CD28/Fc | 65.3 ± 9.1 | 139.8 ± 4.1 |
| CTLA-4/Fc | 41.1 ± 12.0* | 84.9 ± 17.0* |
| ICOS/Fc | 72.5 ± 8.0 | N.T. |
| B7-1(CD80)/Fc | 96.9 ± 13.9 | N.T. |
| B7-H2(B7h)/Fc | 43.7 ± 5.9* | 95.9 ± 13.4* |
| CD28/Fc + B7-H2/Fc | N.T. | 79.5 ± 15.6* |
| CTLA-4/Fc + B7-H2/Fc | N.T. | 71.0 ± 5.0* |

PHA-stimulated CD4+ PBMC were infected with HIV-1 NL4-3, and cultured in the medium containing IL-2 in the presence of 10 µg/ml of the indicated recombinant chimeric proteins of various B7-CD28 family molecules and the human Fc region of IgG. HIV-1 p24 in the supernatant was measured by ELISA after 4 days. The values indicate the mean ± SD from triplicate determinations.

N.T., not tested.

* $P < 0.05$.

following infection with HIV-1 NL4-3. The results are shown in Table 3. B7-H2 (B7h)/Fc and CTLA-4/Fc significantly suppressed HIV-1 replication. CD28/Fc produced low levels of suppression, but the effects of ICOS/Fc were minimal. B7-1 (CD80)/Fc enhanced viral replication. A mixture of B7-H2 (B7h)/Fc and either CD28/Fc or CTLA-4/Fc resulted in maximal suppression. These results indicate that B7-H2 (B7h), a natural ligand of ICOS, suppressed HIV-1 replication in a similar manner to ICOS-mAbs, whereas B7-1, a natural ligand of CD28 and CTLA-4, enhanced HIV-1 replication.

Discussion

In the present study, we showed that ICOS transmitted signals suppressing HIV-1 replication in CD4+ T cells in vitro. ICOS-mAb suppressed HIV-1 replication when the mAb was either added directly to the cell culture or was immobilized. A soluble form of B7-H2 (B7h), a natural ligand for ICOS, also suppressed HIV-1 replication. Immobilized CD28-mAb suppressed HIV-1 replication in CD4+ cells as previously reported (Levine et al., 1996), while direct application of CD28 mAb or its natural ligand B7-1 mostly enhanced viral replication consistently with other reports, indicating that CD28 costimulation supports HIV-1 replication (Diegel et al., 1993).

Immobilized ICOS or CD28 mAbs significantly inhibited replication of both X4 and R5 wild-type HIV-1 (Fig. 2). In addition, these mAbs inhibited the expression of pseudotype HIV-1 possessing HIV-1 envelope but not VSV-G envelope (Fig. 5). Replication of pseudotype HIV-1 with VSV-G-envelope was even enhanced especially when the cells were pretreated with CD28 mAb. Moreover, immobilized ICOS or CD28 mAbs significantly inhibited DNA synthesis

during the initial cycle of viral replication in the cells infected with wild-type HIV-1 (Fig. 4). These observations indicated that the entry stages of the viral life cycle were mainly affected by these mAbs. However, CD4 and CXCR4 expression on the cell surface was not affected by stimulation with ICOS and CD28 mAbs, and CCR5 levels were even enhanced by treatment with these mAbs (Fig. 6). Thus, the suppression of HIV-1 replication by these mAbs was not attributable to the down-regulation of HIV-1 receptors or coreceptors.

The cytokines and chemokines potentially modify HIV-1 replication. Stimulation of CD28 induces IL-2, while stimulation of ICOS results in the production of other cytokines (Beier et al., 2000; Coyle et al., 2000; Hutloff et al., 1999). We measured several cytokines and chemokines in two different culture conditions. When the cells were treated with ICOS or CD28 mAbs immediately after PHA stimulation, ICOS and CD28 stimulation enhanced IL-5 and IL-10, but suppressed TNFα and IFNγ production. When the cells were treated with mAbs 4 days after PHA stimulation, the levels of cytokine production were generally low, and alteration of cytokine levels by ICOS and CD28 mAbs was limited. Because HIV-1 expression was suppressed by these mAbs in both culture conditions as well (Figs. 5B and 7A), alteration of cytokines observed here was unlikely to be a major reason for the viral suppression. The precise mechanisms underlying the suppression of HIV-1 replication by ICOS and CD28 mAbs remains to be determined.

The signaling pathways downstream of CD28 and ICOS are not fully understood. Stimulation of the T cell receptor in concert with CD28 induces activation of JNK, P38 MAPK, AP-1, and interleukin-2 (Avraham et al., 1998; Faris et al., 1996; Hehner et al., 2000), which involves activation of Vav and Rac-1 (Raab et al., 2001; Salojin et al., 1999). Activation of NF-κB also occurs following CD28 costimulation, which results from IκB degradation (Harhaj and Sun, 1998; Kane et al., 2002). Recent reports have shown that MAPKKK might activate the IKK complex, and lead to degradation of IκB following CD28 costimulation (Tuosto et al., 2000). In the present study, ICOS- and CD28-mediated HIV-1 suppression was reversed by the NF-κB inhibitor Leflunomide, which prevents degradation of IκB (Manna and Aggarwal, 1999). Because NF-κB also positively regulates HIV-1 replication itself, Leflunomide alone suppressed HIV-1 production. Nevertheless, the levels of HIV-1 replication in the presence of Leflunomide recovered by stimulation with CD28 mAb to the control levels and also with ICOS mAb to a lesser extent (Fig. 8). Leflunomide has been shown to inhibit protein tyrosine kinase and MAP kinase pathways involving MEK (Brazelton and Morris, 1996; Manna and Aggarwal, 1999). In the presence of PD98059, the MEK inhibitor, CD28, but not ICOS mAb, enhanced HIV-1 replication (Fig. 8). These results implies that the CD28-mediated positive effects on HIV-1 replication overcame its inhibitory effects in the presence of Leflunomide or PD98059, indicating that the CD28-medi-

ated signals suppressing HIV-1 replication might involve the NF- κ B and MEK pathways. ICOS also showed a weak positive effect on HIV-1 replication in the presence of Leflunomide, indicating that its inhibitory effects might involve partly common pathways to those of CD28.

Among the chimeric Fc-fusion proteins with various B7-CD28 family molecules tested, CTLA-4/Fc and B7-H2 (B7h)/Fc significantly inhibited HIV-1 replication in CD4⁺ PBMC (Table 2). CTLA-4/Fc, known to efficiently block interactions between B7-1 (CD80)/B7-2 (CD86) and CD28, suppressed replication of HIV-1. A similar suppression of HIV-1 replication by CTLA-4/Fc has been previously reported (Diegel et al., 1993; Smithgall et al., 1995). This occurs probably because CTLA-4/Fc blocked the positive regulation through B7-1 (CD80)/B7-2 (CD86) that is expressed in a small proportion of PBMC. Although immobilized CD28-mAb mediated HIV-1 suppression *in vitro*, natural ligands of CD28 may preferentially mediate positive signals for HIV-1 replication. This notion is partly supported by previous reports showing that stimulation of CD28 with cross-linking mAb, but not with natural ligand B7-1, could activate ERK2 and p21ras (Nunes et al., 1994).

Because B7-H2 (B7h) is a natural ligand of ICOS, it was not unexpected that B7-H2 (B7h)/Fc suppressed HIV-1, as B7-H2 (B7h)/Fc binds ICOS and potentially activates signals downstream to ICOS on CD4⁺ T cells. This is consistent with our current findings that ICOS mAbs mediated suppression of HIV-1. The simple addition of ICOS/Fc, which is capable of binding B7-H2 (B7h), to the PBMC culture did not alter HIV-1 replication, suggesting that the levels of natural ligands of ICOS present in the culture is insufficient to suppress HIV-1. Nevertheless, suppression of HIV-1 replication by B7-H2 (B7h)/Fc indicates that natural or artificial ligands of ICOS potentially elicit suppressive effects on HIV-1 replication.

B7-H2 (B7h) is more broadly expressed than B7-1 (CD80)/B7-2 (CD86) in both lymphoid and a variety of nonlymphoid tissues (Swallow et al., 1999), implying that ICOS-expressing cells could be regulated by various peripheral cells. A recent clinical study on HIV-1 infection showed that ICOS expression peaked during the early stages of primary infection, dropped to control levels in the asymptomatic period, and subsequently increased as AIDS developed (Lucia et al., 2000). They also showed that HIV-1 gp120 induced transient ICOS expression in naïve T cells. Our results suggest that ICOS stimulation may render uninfected T cells resistant to HIV-1 replication, and less efficiently suppress HIV-1 expression in persistently infected cells. ICOS-mediated suppression of HIV-1 replication may play some roles in regulating HIV-1 *in vivo*.

In conclusion, stimulation of ICOS by mAb or its ligands mediated suppression of the early steps of HIV-1 replication in CD4⁺ PBMC. These findings contribute to our understanding of HIV-1 regulation *in vivo*, and the development of strategies to control HIV-1.

Materials and methods

Cells

PBMC from healthy individuals were isolated by Ficoll-Hypaque density centrifugation. CD4⁺ T cell fractions were negatively purified from PBMC by magnetic cell sorting following incubation with a cocktail of mAbs for CD8, CD14, CD16, CD19, CD36, CD56, CD123, TCR $\gamma\delta$, and Glycophorin A, using MACS Cell Isolation Kits (Miltenyi Biotec, Bergisch Gladbach, Germany). Flow cytometry showed that these fractions contained 93% CD4⁺ cells. Unfractionated PBMC or CD4⁺ T cells were stimulated with 0.05% PHA P (Becton Dickinson, MD) for 30 min, washed, and then cultured in RPMI 1640 (GIBCO BRL, Long Island, NY) supplemented with 10% heat-inactivated fetal calf serum (FCS), 10 U/ml recombinant human interleukin-2 (IL-2) (Shionogi, Osaka, Japan) and 2 mM L-glutamine, 100 U/ml penicillin, and 100 μ g/ml streptomycin sulfate at 37 °C in a 5% CO₂ incubator at a concentration of 1×10^6 cells/ml for 4–6 days. Cells were then used for HIV-1 infection. In some experiments, Dynabeads (Dynal A.S., Oslo, Norway) coated with anti-CD8 mAbs were used to enrich CD4⁺ PBMC. 293T cells were maintained in Dulbecco's Modification of Eagle's Medium (ICN Biomedicals Inc., OH) supplemented with 10% FCS.

Flow cytometric analysis

The cells were incubated with mAbs antigens or control mouse immunoglobulin, washed, and subsequently stained with FITC-conjugated goat anti-mouse IgG + IgM mAbs (Immunotech, Marseilles, France) for 30 min at room temperature. The cells were washed and analyzed using a flow cytometer (FACS Calibur, Becton Dickinson, San Jose, CA).

Antibodies and reagents

The mAbs to human CD28 family molecules used were TN228, anti-CD28 (Lanier et al., 1995), MIH8, anti-CTLA-4, SA12, anti-ICOS (Sakamoto et al., 2001), and MIH4, anti-PD-1 (Youngnak et al., 2003), all of which were prepared as purified mouse IgG1. MIH8 anti-CTLA4 was generated by immunizing DBA/2 mice with human CTLA-4-transfected P815 cells (Oki et al., 1999). SA12 was kindly provided by Japan Tobacco Inc. (Osaka, Japan). Fluorescein isothiocyanate (FITC)-conjugated CD4, phycoerythrin (PE)-conjugated CXCR4 and CCR5, and their isotype-matched control antibodies were purchased from BD Bioscience (San Diego, CA). Human CD28/Fc, CTLA-4/Fc, B7-1/Fc, and B7-H2/Fc, which were chimeric recombinant proteins fused to the Fc region of human IgG, were purchased from R & D Systems, Inc. (Minneapolis, MN). Human ICOS/Fc was provided by Japan Tobacco Inc. A NF- κ B inhibitor Leflunomide, MEK (MAPKK) inhibitor PD98059 (Sigma-

Aldrich, St. Louis, MO), and p38 MAP kinase inhibitor SB202190 (Calbiochem, San Diego, CA) were also used.

Proliferation assay

Cell proliferation was measured using the [^3H]thymidine uptake method. PHA-stimulated CD4 $^+$ T cells cultured for 4–6 days in a medium containing IL-2 were incubated in 96-well flat-bottom plates at a concentration of 10^5 cells/200 μl in either the presence or absence of mAbs for 4 days. [^3H]thymidine (37 kBq) was added into each well during the last 16 h. Cells were then harvested on a glass filter, and the incorporation of [^3H]thymidine into the cells was monitored with a MicroBeta scintillation counting system (Pharmacia Biotech, Cambridge, UK).

Cytotoxicity test

PHA-stimulated CD4 $^+$ T cells cultured in IL-2-containing medium for 4–6 days were incubated in 96-well flat bottom plates at a concentration of 10^5 cells/well in either the presence or absence of mAbs, and LDH released into the supernatant was monitored every 24 h over 3 days using a Non-Radioactive Cytotoxicity Assay Kit (Promega, Madison, WI).

Virus preparation and infection

X4 HIV 1 strain NL4-3 (Adachi et al., 1986) and R5 HIV-1 strain JR-CSF (Koyanagi et al., 1987), kindly provided by Dr. Y. Koyanagi (Tohoku University), were propagated in PBMC and the supernatant was used as virus source. For HIV-1 infection, PHA-stimulated unfractionated or CD4 $^+$ PBMC were cultured in a medium containing IL-2 for 4–6 days, incubated with HIV-1-containing supernatant at a concentration of 4×10^6 /ml for 2 h at 37 °C, washed, and then cultured in a fresh medium containing IL-2. HIV-1 pseudotype virus was prepared as described elsewhere (Masuda et al., 1995; Planelles et al., 1995). Briefly, using Lipofectamine (GIBCO BRL), 10^6 of 293T cells were transfected with 1 μg each of pNL4-3luc Δenv vector and pLET-LAI (Poon et al., 1998) or pHCMVG (Yee et al., 1994), all of which were kindly provided by Dr. Irvin S.Y. Chen, University of California. pNL4-3 luc Δenv is an envelope defective pNL4-3 vector containing the luciferase gene inserted at the *Nef* site (Masuda et al., 1995). pLET-LAI encodes envelope proteins of HIV-1 LAI strain (Poon et al., 1998), and pHCMVG encodes a vesicular stomatitis virus envelope (Yee et al., 1994). Culture supernatant (5 ml) of the transfected 293T cells was harvested 48 h posttransfection, filtered through a 0.45- μm filter, and used as pseudotype HIV-1. For infection, PHA-stimulated CD4 $^+$ T cells were incubated for 3 h with a pseudotype HIV-1-containing supernatant that had been treated with DNase I (140 u/ml, Worthington, NJ) in the presence of 10 mM MgCl_2 at 37 °C for 1 h to avoid

DNA contamination before infection. The cells were then washed and cultured in a IL-2-containing medium at a concentration of 5×10^5 cells/ml in 24- or 12-well plates for 48 h in either the presence or absence of mAbs. Virus replication was determined from luciferase activity.

Treatment of HIV-1-infected cells with mAbs

MABs at concentrations ranging between 0.025 and 10 $\mu\text{g}/\text{ml}$ were either added directly to the culture or pre-coated in the wells before culture. For pre-coating, mAbs diluted in PBS was incubated overnight in 96-well or 48-well flat-bottomed plates at 4 °C. The wells were washed with PBS immediately before the culture of HIV-1-infected cells.

Analysis of HIV-1 cDNA synthesis

HIV-1-infected cells were harvested after various incubation periods, washed, and 10^6 cells were disrupted in a 150 μl urea lysis buffer (4.7 M urea, 1.3% sodium dodecyl sulfate, 0.23 M NaCl, 0.67 mM EDTA [pH 8.0], 6.7 mM Tris-HCl [pH 8.0]) and underwent phenol–chloroform extraction and ethanol precipitation. Precipitated DNA was then resuspended in 30 μl water. A 0.5 μg DNA pellet of each sample underwent PCR using primer pairs (Zack et al., 1990) amplifying a DNA fragment between HIV-1 LTR U3 and HIV gag (M667/M661) for 35 cycles at 95 °C for 1 min, 65 °C for 2 min, and 72 °C for 2 min as previously described (Tsurutani et al., 2000; Zack et al., 1990). For HIV-1 DNA standards, 250–5000 copies of linearized HIV-1 JR-CSF DNA were amplified in parallel. Amplified products were resolved on a 2% agarose gel and stained with SYBER-Green (FMC Bioproduct, Rocklan, MN).

Enzyme-linked immunosorbent assay (ELISA)

HIV-1 p24, human RANTES, IFN- γ , SDF-1 α , and MIP-1 α in cell-free supernatants were measured by HIV-1 p24 Antigen ELISA kit (ZeptoMetrix, NY), and human RANTES, IFN- γ , SDF-1 α , and MIP-1 α immunoassay kit (R & D Systems), respectively. Absorbances were measured at 450 nm using microplate reader (Bio-Rad, Hercules, CA) and analyzed with Microplate Manager III software.

Measurement of cytokines

Human IL-4, IL-5, IL-10, and TNF α in the culture supernatants were measured by the Cytometric Bead Array Kit (BD Biosciences, Pharmingen, CA) in accordance with the manufacturer's instructions.

Luciferase assay

For luciferase analysis, cells were lysed by 100 μl of $1 \times$ luciferase lysis buffer (Promega) 24 h postinfection, and

luciferase expression was measured with a luciferase assay system (Lumat, EG&G Berthold, Germany).

Statistical analysis

Differences among HIV-1 p24 values of individual sample groups vs. controls were evaluated by Dunnett's *t* test using SPSS Base 11.0J (SPSS Inc., Chicago, IL). *P* values less than 0.05 were considered to be statistically significant.

Acknowledgments

We thank Japan Tobacco Inc. (Osaka, Japan) for providing SA12 mAb. This work was supported by grants from the Ministry of Health, Labor, and Welfare of Japan.

References

- Adachi, A., Gendelman, H.E., Koenig, S., Folks, T., Willey, R., Rabson, A., Martin, M.A., 1986. Production of acquired immunodeficiency syndrome-associated retrovirus in human and nonhuman cells transfected with an infectious molecular clone. *J. Virol.* 59 (2), 284–291.
- Avraham, A., Jung, S., Samuels, Y., Seger, R., Ben-Neriah, Y., 1998. Costimulation-dependent activation of a JNK-kinase in T lymphocytes. *Eur. J. Immunol.* 28 (8), 2320–2330.
- Azuma, M., Ito, D., Yagita, H., Okumura, K., Phillips, J.H., Lanier, L.L., Somoza, C., 1993. B70 antigen is a second ligand for CTLA-4 and CD28. *Nature* 366 (6450), 76–79.
- Beier, K.C., Hutloff, A., Dittrich, A.M., Heuck, C., Rauch, A., Buchner, K., Ludewig, B., Ochs, H.D., Mages, H.W., Kroczeck, R.A., 2000. Induction, binding specificity and function of human ICOS. *Eur. J. Immunol.* 30 (12), 3707–3717.
- Brazelton, T.R., Morris, R.E., 1996. Molecular mechanisms of action of new xenobiotic immunosuppressive drugs: tacrolimus (FK506), sirolimus (rapamycin), mycophenolate mofetil and leflunomide. *Curr. Opin. Immunol.* 8 (5), 710–720.
- Brunet, J.F., Denizot, F., Luciani, M.F., Roux-Dosseto, M., Suzan, M., Mattei, M.G., Golstein, P., 1987. A new member of the immunoglobulin superfamily—CTLA-4. *Nature* 328 (6127), 267–270.
- Buonfiglio, D., Bragardo, M., Redoglia, V., Vaschetto, R., Bottarel, F., Bonissoni, S., Bensi, T., Mezzatesta, C., Janeway Jr., C.A., Dianzani, U., 2000. The T cell activation molecule H4 and the CD28-like molecule ICOS are identical. *Eur. J. Immunol.* 30 (12), 3463–3467.
- Coyle, A.J., Lehar, S., Lloyd, C., Tian, J., Delaney, T., Manning, S., Nguyen, T., Burwell, T., Schneider, H., Gonzalo, J.A., Gosselin, M., Owen, L.R., Rudd, C.E., Gutierrez-Ramos, J.C., 2000. The CD28-related molecule ICOS is required for effective T cell-dependent immune responses. *Immunity* 13 (1), 95–105.
- Diegel, M.L., Moran, P.A., Gilliland, L.K., Damle, N.K., Hayden, M.S., Zarling, J.M., Ledbetter, J.A., 1993. Regulation of HIV production by blood mononuclear cells from HIV-infected donors: II. HIV-1 production depends on T cell–monocyte interaction. *AIDS Res. Hum. Retroviruses* 9 (5), 465–473.
- Dong, H., Zhu, G., Tamada, K., Chen, L., 1999. B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion. *Nat. Med.* 5 (12), 1365–1369.
- Faris, M., Kokot, N., Lee, L., Nel, A.E., 1996. Regulation of interleukin-2 transcription by inducible stable expression of dominant negative and dominant active mitogen-activated protein kinase kinase in Jurkat T cells. Evidence for the importance of Ras in a pathway that is controlled by dual receptor stimulation. *J. Biol. Chem.* 271 (44), 27366–27373.
- Finzi, D., Hermankova, M., Pierson, T., Carruth, L.M., Buck, C., Chaisson, R.E., Quinn, T.C., Chadwick, K., Margolick, J., Brookmeyer, R., Gallant, J., Markowitz, M., Ho, D.D., Richman, D.D., Siliciano, R.F., 1997. Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science* 278 (5341), 1295–1300.
- Frauwirth, K.A., Thompson, C.B., 2002. Activation and inhibition of lymphocytes by costimulation. *J. Clin. Invest.* 109 (3), 295–299.
- Freeman, G.J., Long, A.J., Iwai, Y., Bourque, K., Chernova, T., Nishimura, H., Fitz, L.J., Malenkovich, N., Okazaki, T., Byrne, M.C., Horton, H.F., Fouser, L., Carter, L., Ling, V., Bowman, M.R., Carreno, B.M., Collins, M., Wood, C.R., Honjo, T., 2000. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J. Exp. Med.* 192 (7), 1027–1034.
- Hara, T., Fu, S.M., Hansen, J.A., 1985. Human T cell activation. II. A new activation pathway used by a major T cell population via a disulfide-bonded dimer of a 44 kilodalton polypeptide (9.3 antigen). *J. Exp. Med.* 161 (6), 1513–1524.
- Harhaj, E.W., Sun, S.C., 1998. IkappaB kinases serve as a target of CD28 signaling. *J. Biol. Chem.* 273 (39), 25185–25190.
- Hehner, S.P., Hofmann, T.G., Dienz, O., Droge, W., Schmitz, M.L., 2000. Tyrosine-phosphorylated Vav1 as a point of integration for T-cell receptor- and CD28-mediated activation of JNK, p38, and interleukin-2 transcription. *J. Biol. Chem.* 275 (24), 18160–18171.
- Ho, D.D., Moudgil, T., Alam, M., 1989. Quantitation of human immunodeficiency virus type 1 in the blood of infected persons. *N. Engl. J. Med.* 321 (24), 1621–1625.
- Ho, D.D., Neumann, A.U., Perelson, A.S., Chen, W., Leonard, J.M., Markowitz, M., 1995. Rapid turnover of plasma virions and CD4 lymphocytes in HIV-1 infection. *Nature* 373 (6510), 123–126.
- Hutloff, A., Dittrich, A.M., Beier, K.C., Eljaschewitsch, B., Kraft, R., Anagnostopoulos, I., Kroczeck, R.A., 1999. ICOS is an inducible T-cell co-stimulator structurally and functionally related to CD28. *Nature* 397 (6716), 263–266.
- Kane, L.P., Lin, J., Weiss, A., 2002. It's all Rel-ative: NF-kappaB and CD28 costimulation of T-cell activation. *Trends Immunol.* 23 (8), 413–420.
- Kannagi, M., Masuda, T., Hattori, T., Kanoh, T., Nasu, K., Yamamoto, N., Harada, S., 1990. Interference with human immunodeficiency virus (HIV) replication by CD8+ T cells in peripheral blood leukocytes of asymptomatic HIV carriers in vitro. *J. Virol.* 64 (7), 3399–3406.
- Koyanagi, Y., Miles, S., Mitsuyasu, R.T., Merrill, J.E., Vinters, H.V., Chen, I.S., 1987. Dual infection of the central nervous system by AIDS viruses with distinct cellular tropisms. *Science* 236 (4803), 819–822.
- Krummel, M.F., Allison, J.P., 1995. CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation. *J. Exp. Med.* 182 (2), 459–465.
- Lanier, L.L., O'Fallon, S., Somoza, C., Phillips, J.H., Linsley, P.S., Okumura, K., Ito, D., Azuma, M., 1995. CD80 (B7) and CD86 (B70) provide similar costimulatory signals for T cell proliferation, cytokine production, and generation of CTL. *J. Immunol.* 154 (1), 97–105.
- Latchman, Y., Wood, C.R., Chernova, T., Chaudhary, D., Borde, M., Chernova, I., Iwai, Y., Long, A.J., Brown, J.A., Nunes, R., Greenfield, E.A., Bourque, K., Boussiotis, V.A., Carter, L.L., Carreno, B.M., Malenkovich, N., Nishimura, H., Okazaki, T., Honjo, T., Sharpe, A.H., Freeman, G.J., 2001. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat. Immunol.* 2 (3), 261–268.
- Lenschow, D.J., Walunas, T.L., Bluestone, J.A., 1996. CD28/B7 system of T cell costimulation. *Annu. Rev. Immunol.* 14, 233–258.
- Levine, B.L., Mosca, J.D., Riley, J.L., Carroll, R.G., Vahey, M.T., Jagodzinski, L.L., Wagner, K.F., Mayers, D.L., Burke, D.S., Weislow, O.S., St Louis, D.C., June, C.H., 1996. Antiviral effect and ex vivo CD4+ T cell proliferation in HIV-positive patients as a result of CD28 costimulation. *Science* 272 (5270), 1939–1943.
- Linsley, P.S., Clark, E.A., Ledbetter, J.A., 1990. T-cell antigen CD28 medi-

- ates adhesion with B cells by interacting with activation antigen B7/BB-1. *Proc. Natl. Acad. Sci. U. S. A.* 87 (13), 5031–5035.
- Lucia, M.B., Buonfiglio, D., Bottarel, F., Bensi, T., Rutella, S., Rumi, C., Ortona, L., Janeway Jr., C.A., Cauda, R., Dianzani, U., 2000. Expression of the novel T cell activation molecule hpH4 in HIV-infected patients: correlation with disease status. *AIDS Res. Hum. Retroviruses* 16 (6), 549–557.
- Manna, S.K., Aggarwal, B.B., 1999. Immunosuppressive leflunomide metabolite (A77 1726) blocks TNF-dependent nuclear factor-kappa B activation and gene expression. *J. Immunol.* 162 (4), 2095–2102.
- Masuda, T., Planelles, V., Krogstad, P., Chen, I.S., 1995. Genetic analysis of human immunodeficiency virus type 1 integrase and the U3 att site: unusual phenotype of mutants in the zinc finger-like domain. *J. Virol.* 69 (11), 6687–6696.
- Nishimura, H., Honjo, T., 2001. PD-1: an inhibitory immunoreceptor involved in peripheral tolerance. *Trends Immunol.* 22 (5), 265–268.
- Nunes, J.A., Collette, Y., Truneh, A., Olive, D., Cantrell, D.A., 1994. The role of p21ras in CD28 signal transduction: triggering of CD28 with antibodies, but not the ligand B7-1, activates p21ras. *J. Exp. Med.* 180 (3), 1067–1076.
- Oki, S., Kohsaka, T., Azuma, M., 1999. Augmentation of CTLA-4 expression by wortmannin: involvement of lysosomal sorting properties of CTLA-4. *Int. Immunol.* 11 (9), 1563–1571.
- Pantaleo, G., Graziosi, C., Demarest, J.F., Butini, L., Montroni, M., Fox, C.H., Orenstein, J.M., Kotler, D.P., Fauci, A.S., 1993. HIV infection is active and progressive in lymphoid tissue during the clinically latent stage of disease. *Nature* 362 (6418), 355–358.
- Perelson, A.S., Neumann, A.U., Markowitz, M., Leonard, J.M., Ho, D.D., 1996. HIV-1 dynamics in vivo: virion clearance rate, infected cell lifespan, and viral generation time. *Science* 271 (5255), 1582–1586.
- Piatak Jr., M., Saag, M.S., Yang, L.C., Clark, S.J., Kappes, J.C., Luk, K.C., Hahn, B.H., Shaw, G.M., Lifson, J.D., 1993. High levels of HIV-1 in plasma during all stages of infection determined by competitive PCR. *Science* 259 (5102), 1749–1754.
- Planelles, V., Haislip, A., Withers-Ward, E.S., Stewart, S.A., Xie, Y., Shah, N.P., Chen, I.S., 1995. A new reporter system for detection of retroviral infection. *Gene Ther.* 2 (6), 369–376.
- Poon, B., Grovit-Ferbas, K., Stewart, S.A., Chen, I.S., 1998. Cell cycle arrest by Vpr in HIV-1 virions and insensitivity to antiretroviral agents. *Science* 281 (5374), 266–269.
- Raab, M., Pfister, S., Rudd, C.E., 2001. CD28 signaling via VAV/SLP 76 adaptors: regulation of cytokine transcription independent of TCR ligation. *Immunity* 15 (6), 921–933.
- Sakamoto, S., Tezuka, K., Tsuji, T., Hori, N., Tamatani, T., 2001. AILIM/ICOS: its expression and functional analysis with monoclonal antibodies. *Hybrid. Hybridomics* 20 (5–6), 293–303.
- Salojin, K.V., Zhang, J., Delovitch, T.L., 1999. TCR and CD28 are coupled via ZAP-70 to the activation of the Vav/Rac-1/PAK-1/p38 MAPK signaling pathway. *J. Immunol.* 163 (2), 844–853.
- Sharpe, A.H., Freeman, G.J., 2002. The B7-CD28 superfamily. *Nat. Rev. Immunol.* 2 (2), 116–126.
- Smithgall, M.D., Wong, J.G., Linsley, P.S., Haffar, O.K., 1995. Costimulation of CD4+ T cells via CD28 modulates human immunodeficiency virus type 1 infection and replication in vitro. *AIDS Res. Hum. Retroviruses* 11 (8), 885–892.
- Swallow, M.M., Wallin, J.J., Sha, W.C., 1999. B7h, a novel costimulatory homolog of B7.1 and B7.2, is induced by TNFalpha. *Immunity* 11 (4), 423–432.
- Tamatani, T., Tezuka, K., Hanzawa-Higuchi, N., 2000. AILIM/ICOS: a novel lymphocyte adhesion molecule. *Int. Immunol.* 12 (1), 51–55.
- Tseng, S.Y., Otsuji, M., Gorski, K., Huang, X., Slansky, J.E., Pai, S.I., Shalabi, A., Shin, T., Pardoll, D.M., Tsuchiya, H., 2001. B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells. *J. Exp. Med.* 193 (7), 839–846.
- Tsurutani, N., Kubo, M., Maeda, Y., Ohashi, T., Yamamoto, N., Kannagi, M., Masuda, T., 2000. Identification of critical amino acid residues in human immunodeficiency virus type 1 IN required for efficient proviral DNA formation at steps prior to integration in dividing and nondividing cells. *J. Virol.* 74 (10), 4795–4806.
- Tuosto, L., Costanzo, A., Guido, F., Marinari, B., Vossio, S., Moretti, F., Levrero, M., Piccolella, E., 2000. Mitogen-activated kinase kinase 1 regulates T cell receptor- and CD28-mediated signaling events which lead to NF-kappaB activation. *Eur. J. Immunol.* 30 (9), 2445–2454.
- Walker, C.M., Moody, D.J., Stites, D.P., Levy, J.A., 1986. CD8+ lymphocytes can control HIV infection in vitro by suppressing virus replication. *Science* 234 (4783), 1563–1566.
- Walker, B.D., Chakrabarti, S., Moss, B., Paradis, T.J., Flynn, T., Durno, A.G., Blumberg, R.S., Kaplan, J.C., Hirsch, M.S., Schooley, R.T., 1987. HIV-specific cytotoxic T lymphocytes in seropositive individuals. *Nature* 328 (6128), 345–348.
- Walunas, T.L., Lenschow, D.J., Bakker, C.Y., Linsley, P.S., Freeman, G.J., Green, J.M., Thompson, C.B., Bluestone, J.A., 1994. CTLA-4 can function as a negative regulator of T cell activation. *Immunity* 1 (5), 405–413.
- Wei, X., Ghosh, S.K., Taylor, M.E., Johnson, V.A., Emini, E.A., Deutsch, P., Lifson, J.D., Bonhoeffer, S., Nowak, M.A., Hahn, B.H., et al., 1995. Viral dynamics in human immunodeficiency virus type 1 infection. *Nature* 373 (6510), 117–122.
- Wong, J.K., Hezareh, M., Gunthard, H.F., Havlir, D.V., Ignacio, C.C., Spina, C.A., Richman, D.D., 1997. Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. *Science* 278 (5341), 1291–1295.
- Yee, J.K., Friedmann, T., Burns, J.C., 1994. Generation of high-titer pseudotyped retroviral vectors with very broad host range. *Methods Cell Biol.* 43 (Pt. A), 99–112.
- Yoshinaga, S.K., Whoriskey, J.S., Khare, S.D., Sarmiento, U., Guo, J., Horan, T., Shih, G., Zhang, M., Coccia, M.A., Kohno, T., Tafuri-Bladt, A., Brankow, D., Campbell, P., Chang, D., Chiu, L., Dai, T., Duncan, G., Elliott, G.S., Hui, A., McCabe, S.M., Scully, S., Shahinian, A., Shaklee, C.L., Van, G., Mak, T.W., et al., 1999. T-cell co-stimulation through B7RP-1 and ICOS. *Nature* 402 (6763), 827–832.
- Younghak, P., Kozono, Y., Kozono, H., Iwai, H., Otsuki, N., Jin, H., Omura, K., Yagita, H., Pardoll, D.M., Chen, L., Azuma, M., 2003. Differential binding properties of B7-H1 and B7-DC to programmed death-1. *Biochem. Biophys. Res. Commun.* 307 (3), 672–677.
- Zack, J.A., Arrigo, S.J., Weitsman, S.R., Go, A.S., Haislip, A., Chen, I.S., 1990. HIV-1 entry into quiescent primary lymphocytes: molecular analysis reveals a labile, latent viral structure. *Cell* 61 (2), 213–222.